

14 Other organelles of mitochondrial origin — 73

15 Genomes are not alive — 78

Part II: Well-studied examples

16 Anaerobic use of the mitochondrial electron-transport chain — 85

17 *Naegleria gruberi*, a strict aerobe with an “anaerobic genome” — 89

18 Malate dismutation in the liver fluke *Fasciola hepatica* — 92

19 The roundworms *Ascaris suum* and *Ascaris lumbricoides* — 97

20 Animals in tidal zones, anaerobic sediments and sulfide — 101

20.1 The mollusc *Mytilus edulis* — 103

20.2 The polychaete annelid *Arenicola marina* — 105

20.3 The peanut worm *Sipunculus nudus* — 110

20.4 Diverse physiological functions of H₂S in animals — 112

21 Anaerobic respiration in eukaryotes, rare but there — 115

21.1 Nitrate respiration in *Fusarium* and *Cylindrocarpon* — 119

22 Enzymes of anaerobic energy metabolism in algae — 124

23 Wax ester fermentation in *Euglena gracilis* — 127

24 *Chlamydomonas reinhardtii*, a jack of all trades — 131

25 Organisms with hydrogenosomes — 136

25.1 Discovery — 136

25.2 Trichomonads — 139

25.3 Chytrids — 145

25.4 Ciliates — 150

26 *Nyctotherus ovalis* and H₂-producing mitochondria — 151

26.1 *Blastocystis hominis* — 155