



# Regulations for calcium, magnesium or hardness in drinking water in the European Union member states



Frantisek Kozisek

National Institute of Public Health, Srobarova 48, Prague, CZ, 10000, Czech Republic

## ARTICLE INFO

### Keywords:

Drinking water  
Calcium  
Magnesium  
Water hardness  
Regulation

## ABSTRACT

Very low or very high concentrations of calcium (Ca) and magnesium (Mg) or total hardness in drinking water have been empirically recognized as the cause of the problems with corrosion, scaling, or taste of water. A large body of scientific evidence over the last 60 years has also attributed health problems to both extremes of these minerals in water. For example, drinking water low in Mg significantly increases the likelihood of cardiovascular mortality. This knowledge is not properly considered in the drinking water quality regulations, especially as more and more water is now produced by desalination disturbing its natural mineral composition. Although the current EU Drinking Water Directive does not include limits for hardness, Ca or Mg, most member states regulate these parameters either through law (12 countries) or technical guidance (7 countries), and several countries have issued educational information for the public. However, the regulations in most countries are not based on current research and not comparable because some provide a recommended range while others provide mandatory minimum or maximum limit values. This summary of the situation in 28 European Union countries is intended to provide information that can contribute to the implementation of the revised EU Directive.

## 1. Introduction

Drinking water is not and cannot be pure H<sub>2</sub>O like distilled or osmotic water. The presence of certain amounts of total dissolved solids (TDS) and some essential elements are needed not only to prevent pipe corrosion and ensure an acceptable taste but also to prevent subacute and chronic adverse health effects from the long-term consumption of water (Rosborg and Kozisek, 2020). Thus, the focus on current microbiological and toxicological limits is inadequate and should be extended to include certain minerals.

Subacute health effects, which may appear within weeks to months after starting regular consumption of distilled or demineralized water, comprise signs of profound deficiency of calcium, magnesium or sodium: extreme fatigue, malaise, nausea, headache, brittleness of nails and hairs, leg and abdominal cramps, preeclampsia, twitch, metabolic acidosis, cardiovascular disorders (arrhythmia), or higher diuresis (Kozisek, 2005). These symptoms are mostly independent of the nutritional status of ill person.

Epidemiologic studies completed in different countries by different teams of researchers since 1960 have consistently reported chronic health effects of naturally occurring low mineral water or artificially softened water (specifically water low in magnesium (Mg), calcium (Ca) or bicarbonates). The most comprehensive review of about two

thousand papers, including more than 100 studies with primary data, was ordered by the Drinking Water Inspectorate (England and Wales) and completed by the University of East Anglia in Norwich in 2005 (UoEA and DWI, 2005). A meta-analysis of 14 analytical observational studies (i.e. the most valid epidemiological studies) that investigated the association between cardiovascular disease and drinking water hardness found convincing evidence of the protective role of Mg in drinking water. The pooled odds ratio showed a statistically significant inverse association between Mg and cardiovascular mortality (OR 0.75; 95%CI 0.68, 0.82;  $p = 0.001$ ). The highest exposure category (people consuming drinking water with Mg content of 8.3–19.4 mg/l) was significantly associated with a decreased likelihood of cardiovascular mortality by 25% compared with people consuming water with Mg content of 2.5–8.2 mg/l (Catling et al., 2008). Two recent and independent meta-analyses that considered several more recent studies (Jiang et al., 2016; Gianfredi et al., 2017) reported similar results. Gianfredi et al. also found a statistically significant protective effect of water calcium on cardiovascular diseases. The current position of the World Health Organization (WHO, 2011, 2017a) is that the strength of evidence is insufficient to propose guidelines for Ca, Mg or water hardness. However, in another WHO publication on the public health significance of Ca and Mg, Calderon and Hunter (2009) concluded “There is a growing consensus among epidemiologists that the

E-mail address: [frantisek.kozisek@szu.cz](mailto:frantisek.kozisek@szu.cz).

<https://doi.org/10.1016/j.yrtph.2020.104589>

Received 23 October 2019; Received in revised form 24 January 2020; Accepted 27 January 2020

Available online 30 January 2020

0273-2300/ © 2020 Elsevier Inc. All rights reserved.




Fig. 1. Comparisons of limit values or recommended ranges for drinking water calcium (A), magnesium (B) and hardness (C) in different EU member states. Data include both legal regulations and technical guidelines.

## Disclaimer

The author declares no conflicts of interest.

## Funding body information

The study was supported by Ministry of Health, Czech Republic – conceptual development of research organization (National Institute of Public Health – NIPH, IN 75010330).

Ministry of Health of the Czech Republic, Palackeho namesti 4, CZ-12801 Prague.

It was also partially supported by the project “Improvement of health status of population of the Slovak Republic through drinking water re-carbonization” (LIFE17 ENV/SK/000036), which is financially supported by the EU’s funding instrument (LIFE Environment and Resource Efficiency) and the Ministry of the Environment of the Slovak Republic.

## Declaration of competing interest

The author declares that he has no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgments

The author would like to thank all ENDWARE members to share their national drinking water regulations.

## References

Calderon, R., Hunter, P., 2009. Epidemiological studies and the association of cardiovascular disease risks with water hardness. In: *Calcium and Magnesium in Drinking-Water. Public Health Significance*. WHO, Geneva, pp. 110–144.

Catling, L.A., Abubakar, I., Lake, I.R., Swift, L., Hunter, P.R., 2008. A systematic review of analytical observational studies investigating the association between cardiovascular disease and drinking water hardness. *J. Water Health* 6 (4), 433–442. <https://doi.org/10.2166/wh.2008.054>.

Costi, D., Calcaterra, P.G., Iori, N., Vourna, S., Nappi, G., Passeri, M., 1999. Importance of bioavailable calcium in drinking water for the maintenance of bone mass in post-menopausal women. *J. Endocrinol. Invest.* 22 (11), 852–856. <https://doi.org/10.1007/BF03343658>.

Curhan, G.C., Willett, W.C., Rimm, E.B., Stampfer, M.J., 1993. A prospective study of dietary calcium and other nutrients and the risk of symptomatic kidney stones. *N. Engl. J. Med.* 328 (12), 833–838. <https://doi.org/10.1056/NEJM199303253281203>.

Council of the European Union, 2019. Proposal for a Directive of the European Parliament and of the Council on the quality of water intended for human consumption (recast) – general approach. Brussels, 6876/19 27 Feb. 2019.

Dahl, C., Søgaard, A.J., Tell, G.S., Flaten, T.P., Hongve, D., Omsland, T.K., Holvik, K., Meyer, H.E., Aamodt, G., 2013. Nationwide data on municipal drinking water and hip fracture: could calcium and magnesium be protective? A NOREPOS study. *Bone* 57 (2), 84–91. <https://doi.org/10.1016/j.bone.2013.06.017>.

Dahl, C., Søgaard, A.J., Tell, G.S., Forsén, L., Flaten, T.P., Hongve, D., Omsland, T.K., Holvik, K., Meyer, H.E., Aamodt, G., 2015. Population data on calcium in drinking water and hip fracture: an association may depend on other minerals in water. A NOREPOS study. *Bone* 81, 292–299. <https://doi.org/10.1016/j.bone.2015.07.020>.

EC (European Community), 1998. Council Directive 98/83/EC of 3 November 1998 on the quality of water intended for human consumption. *Off. J. Eur. Communities - Legislation* 330 (5), 32–54. 12.1998. [https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=uriserv:OJ.L\\_.1998.330.01.0032.01 ENG&toc=OJ:L:1998:330:TOC](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=uriserv:OJ.L_.1998.330.01.0032.01 ENG&toc=OJ:L:1998:330:TOC).

EC (European Commission), 2018. Proposal for a Directive of the European Parliament and of the Council on the quality of water intended for human consumption. COM (2017), 753. 1 Feb. 2018. Available on line. [https://eur-lex.europa.eu/resource.html?uri=cellar:8c5065b2-074f-11e8-b8f5-01aa75ed71a1.0016.02/DOC\\_1&format=PDF](https://eur-lex.europa.eu/resource.html?uri=cellar:8c5065b2-074f-11e8-b8f5-01aa75ed71a1.0016.02/DOC_1&format=PDF), Accessed date: 18 September 2019.

EEC (European Economic Community), 1980. Council Directive 80/778/EEC of 15 July 1980 relating to the quality of water intended for human consumption. *Off. J. Eur. Communities - Legislation* 229, 11–29. 30.38.1980. [https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=uriserv:OJ.L\\_.1980.229.01.0011.01 ENG&toc=OJ:L:1980:229:TOC](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=uriserv:OJ.L_.1980.229.01.0011.01 ENG&toc=OJ:L:1980:229:TOC).

Gianfredi, V., Bragazzi, N.L., Nucci, D., Villarini, M., Moretti, M., 2017. Cardiovascular diseases and hard drinking waters: implications from a systematic review with meta-analysis of case-control studies. *J. Water Health* 15 (1), 31–40. <https://doi.org/10.2166/wh.2016.131>.

Huang, Y., Wang, J., Tan, Y., Wang, L., Lin, H., Lan, L., Xiong, Y., Huang, W., Shu, W., 2018. Low-mineral direct drinking water in school may retard height growth and increase dental caries in schoolchildren in China. *Environ. Int.* 115, 104–109. <https://doi.org/10.1016/j.envint.2018.02.021>.

Huang, Y., Ma, X., Tan, Y., Wang, L., Wang, J., Lan, L., Oiu, Z., Luo, J., Zeng, H., Shu, W., 2019. Consumption of very low mineral water is associated with lower bone mineral content in children. *J. Nutr.* 149, 1994–2000. <https://doi.org/10.1093/jn/nxz161>.

Jiang, L., He, P., Chen, J., Liu, Y., Liu, D., Qin, G., Tan, N., 2016. Magnesium levels in drinking water and coronary heart disease mortality risk: a meta-analysis. *Nutrients* 8 (1), 5. <https://doi.org/10.3390/nu8010005>.

Kozisek, F., 2005. Health risks from drinking demineralized water. In: *Nutrients in Drinking Water*. WHO, Geneva, pp. 148–163.

Nriagu, J., Darroudi, F., Shomar, B., 2016. Health effects of desalinated water: role of electrolyte disturbance in cancer development. *Environ. Res.* 150, 191–204. <https://doi.org/10.1016/j.envres.2016.05.038>.

Rosborg, I., Kozisek, F. (Eds.), 2020. *Drinking Water Minerals and Mineral Balance*, second ed. Springer International Publishing, London. <https://doi.org/10.1007/978-3-030-18034-8>.

Schwartz, B.F., Schenkman, N.S., Bruce, J.E., Leslie, S.W., Stoller, M.L., 2002. Calcium nephrolithiasis: effect of water hardness on urinary electrolytes. *Urology* 60 (1), 23–27. [https://doi.org/10.1016/s0090-4295\(02\)01631-x](https://doi.org/10.1016/s0090-4295(02)01631-x).

Shlezinger, M., Amitai, Y., Goldenberg, I., Shechter, M., 2016. Desalinated seawater supply and all-cause mortality in hospitalized acute myocardial infarction patients from the Acute Coronary Syndrome Israeli Survey 2002–2013. *Int. J. Cardiol.* 220, 544–550. <https://doi.org/10.1016/j.ijcard.2016.06.241>.

Shlezinger, M., Amitai, Y., Akiriv, A., Gabay, H., Shechter, M., Leventer-Roberts, M., 2018. Association between exposure to desalinated sea water and ischemic heart disease, diabetes mellitus and colorectal cancer; A population-based study in Israel. *Environ. Res.* 166, 620–627. <https://doi.org/10.1016/j.envres.2018.06.053>.

UoEA (University of East Anglia), DWI (Drinking Water Inspectorate), 2005. Review of evidence for relationship between incidence of cardiovascular disease and water hardness. Final report, Norwich – London. Available on line. <http://dwi.defra.gov>.